

# Ditrac<sup>®</sup> RAT & MOUSE BAIT

## Cereal Bait with Diphacinone

DITRAC Rat and Mouse Bait is a cereal bait containing the anticoagulant Diphacinone. Diphacinone is more acute than coumarin type anticoagulants and kills rats and mice with less chemical than many other anticoagulants.

DITRAC contains only food-grade cereals and enhancers for optimal acceptance because success with diphacinone baits depends on rodent acceptance and repeated feedings. Bait shyness is not a problem, and the safety factor for non-target animals is high with Diphacinone baits due to the need for repeat feedings. Also the antidote, Vitamin K<sub>1</sub>, is readily available.

DITRAC Rat and Mouse Bait is authorized by USDA for use in federally inspected meat and poultry plants.

DITRAC is packaged in bulk and polyglassine place pacs which protect the bait from dust and chemical odors. These slender place pacs readily fit down burrows, in runways, and in bait stations. The 3-oz. pacs are ideal for rats and the 1½-oz. for mice.

**Packaging:** 25 lb. bulk pail – EPA REG. NO. 12455-19  
60 x 3 oz. place pacs – EPA REG. NO. 12455-78  
120 x 1½ oz. place pacs – EPA REG. NO. 12455-29



## Specimen Label

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions.

**IMPORTANT:** Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

**USE RESTRICTIONS:** For control of Norway rats, roof rats, and house mice in and around homes, industrial and agricultural buildings, and similar man-made structures.

Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

## Ditrac<sup>®</sup> RAT & MOUSE BAIT

### KILLS NORWAY RATS, ROOF RATS, AND HOUSE MICE

ACTIVE INGREDIENT: Diphacinone;  
2-(Diphenylacetyl)-1,3-indandione ..... 0.005%  
INERT INGREDIENTS..... 99.995%  
TOTAL: 100.000%

### KEEP OUT OF REACH OF CHILDREN CAUTION

(See right panel for additional precautionary statements.)

For 24-hour emergency medical advice, call 1-877-854-2494.

Manufactured by:



**Bell Laboratories, Inc.**  
Madison, WI 53704 U.S.A.

EPA EST. NO. 12455-WI-1

EPA REG. NO. 12455-19

### DIRECTIONS FOR USE (Continued from other panel)

#### APPLICATION DIRECTIONS:

**RATS:** Apply 3 to 16 oz. of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

**MICE:** Apply ¼ to ½ oz. (1 to 2 level tablespoons) of bait at 8- to 12- foot intervals. Larger placements (up to 2 oz.) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

**RATS AND MICE:** Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store only in original container in a dry place inaccessible to children and pets.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty pail. Then dispose of empty pail in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

#### PRECAUTIONARY STATEMENTS HAZARD TO HUMANS & DOMESTIC ANIMALS CAUTION

Keep away from humans, domestic animals, and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

**NOTE TO PHYSICIAN:** If ingested, administer Vitamin K<sub>1</sub> intramuscularly or orally as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

#### ENVIRONMENTAL HAZARD

Keep out of lakes, streams or ponds.

**WARRANTY:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.